

**ANTI-DIARRHEAL- loperamide hydrochloride capsule, liquid filled  
QUALITY CHOICE (Chain Drug Marketing Association)**

-----

**Drug Facts**

**Active ingredient (in each capsule)**

Loperamide HCl 2 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

- if you have bloody or black stool
- in children under 12 years of age

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease

**Ask a doctor or pharmacist before use if you are**

taking antibiotics.

**When using this product**

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- **not for use in children under 12 years of age**
- adults and children 12 years of age and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours

**Other information**

- store at 20°-25°C (68°-77°F) Protect from light.
- avoid excessive heat above 40°C (104°F)
- do not use if carton or blister unit is open or torn
- see side panel for lot number and expiration date

**Inactive ingredients**

butylated hydroxyanisole, edible ink, FD&C blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water

**Questions or comments?**

Call **1-248-449-9300 Monday-Friday 9AM-5PM EST**

**Principal Display Panel**

†Compare to the active ingredient in Imodium® A-D

**Anti-Diarrheal**

Controls The Symptoms Of Diarrhea

Loperamide HCl, 2 mg

Suitable For Adults And Children 12 Years And Over

\*Each Liquid-Filled Capsule Contains 2 mg Loperamide HCl

Softgels\*

Distributed by C.D.M.A., Inc.®

43157 W. Nine Mile

Novi, MI 48376-0995

[www.qualitychoice.com](http://www.qualitychoice.com)

Questions: 248-449-9300

**THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

†This product is not manufactured or distributed by McNeil Consumer & Specialty Pharmaceuticals, distributor of Imodium® A-D.

**Product Label**

This product is not manufactured or distributed by McNeil Consumer & Specialty Pharmaceuticals, distributor of Imodium® A-D.

<p><b>Questions or comments?</b> Call 1-248-449-9300 Monday-Friday 9AM-5PM EST</p>	<p>◀</p>
<p><b>Inactive ingredients</b> butylated hydroxyanisole, edible ink, FD&amp;C blue #1, gelatin, glycerin, glyceryl caprylate, polyoxy 40 hydrogenated castor oil, purified water</p>	<p><b>Stop use and ask a doctor if</b> ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition.</p>
<p><b>Other information</b> ■ store at 20°-25°C (68°-77°F). Protect from light. ■ avoid excessive heat above 40°C (104°F) ■ do not use if carton or blister unit is open or torn ■ see side panel for lot number and expiration date</p>	<p><b>When using this product</b> tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.</p>
<p><b>Directions</b> ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ not for use in children under 12 years of age ■ adults and children 12 years of age and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours</p>	<p><b>Ask a doctor before use if you have</b> ■ fever ■ mucus in the stool ■ a history of liver disease <b>Do not use</b> ■ if you have bloody or black stool ■ in children under 12 years of age</p>
<p><b>Drug Facts (continued)</b> If pregnant or breast-feeding, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p><b>Warnings</b> <b>Allergy alert:</b> Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. <b>Heart alert:</b> Taking more than directed can cause serious heart problems or death. <b>Use</b> Controls symptoms of diarrhea, including Travelers' Diarrhea. <b>Active ingredient (in each capsule)</b> Loperamide HCl 2 mg.....Anti-diarrheal <b>Purpose</b></p>



# Anti-Diarrheal

Controls Symptoms Of Diarrhea



NDC 63868-748-12

† Compare to the active ingredient in Imodium® A-D

# Anti-Diarrheal

Controls Symptoms Of Diarrhea

TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

SEE WARNINGS AND PRODUCT INFORMATION.

# Loperamide HCl, 2 mg

Suitable For Adults And Children  
12 Years And Over

\*Each Liquid-Filled Capsule  
Contains 2 mg Loperamide HCl



**12** Softgels\*

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND  
KEEP OUTER CARTON FOR COMPLETE

Lot No.:  
Exp. Date:



Distributed by C.D.M.A., Inc.©  
43157 W. Nine Mile  
Novi, MI 48375  
www.qualitychoice.com  
Questions: 248-449-9300

REV # 9/16 PLD-E45J FC003593



**QUALITY CHOICE Anti-Diarrheal**

## ANTI-DIARRHEAL

loperamide hydrochloride capsule, liquid filled

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-748
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII: 6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>GLYCERYL MONOCAPRYLATE</b> (UNII: TM2TZD4G4A)	
<b>POLYOXYL 40 HYDROGENATED CASTOR OIL</b> (UNII: 7YC686GQ8F)	
<b>WATER</b> (UNII: 059QF0KO0R)	

**Product Characteristics**

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	P13
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-748-12	1 in 1 CARTON	05/31/2013	12/29/2023
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021855	05/31/2013	12/29/2023

**Labeler** - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 11/2021

QUALITY CHOICE (Chain Drug Marketing Association)